

Bhattacharyya statistical divergence of quantum observables

V. Majerník*

Institute of Mathematics, Slovak Academy of Sciences, Bratislava, Štefánikova 47, Slovak Republic

S.Shpyrko†

*Department of Optics, Faculty of Science, Palacký University,
Tr. 17. listopadu 50, CZ-77207 Olomouc, Czech Republic and
Institute of Nuclear Research, Ukrainian Academy of Sciences, pr. Nauki 47 Kiev, Ukraine*

In this article we exploit the Bhattacharyya statistical divergence to determine the similarity of probability distributions of quantum observables. After brief review of useful characteristics of the Bhattacharyya divergence we apply it to determine the similarity of probability distributions of two non-commuting observables. An explicit expression for the Bhattacharyya statistical divergence is found for the case of two observables which are the x- and z-components of the angular momentum of a spin-1/2 system. Finally, a note is given of application of the considered statistical divergence to the specific physical measurement.

PACS numbers: 03.65.Ta; 02.50.-r

Keywords: statistical divergence, probability systems, quantum observables

I. INTRODUCTION

One of the important problems in the probability theory is to find an appropriate measure of the difference or the statistical *divergence* of two probability distributions P and P' . This measure quantifies the degree of the similarity between them. In the mathematical statistics the divergence of two probability distributions is introduced as follows: If $[X, P]$ and $[X, P']$ are two probability spaces then the so-called (Csiszár's) f-divergence of probability distributions P and P' is given as

$$D_f(P; P') = \sum_{x \in X} P'(x) f \left[\frac{P(x)}{P'(x)} \right],$$

where $f(u)$ represents a convex function in the interval $(0, \infty)$ and strictly convex for $u = 1$ [1].

Among the existing divergence measures of two discrete probabilities, $P \equiv [p_1, p_2, \dots, p_n]$ and $Q \equiv [q_1, q_2, \dots, q_n]$, the Kullback-Leibler statistical divergence [3]

$$D_K(P : Q) = \sum_{i=1}^n p_i \log \left(\frac{p_i}{q_i} \right),$$

is perhaps best known and most widely used. This is why this measure has several desirable properties, such as nonnegativity and additivity, which are crucial in its applications. $D(P; Q)$ is *not* symmetrical regarding the exchange of P and Q . For $D_K(P; Q)$, the inequality holds

$$\sum_{i=1}^n p_i \log \frac{p_i}{q_i} \geq 0.$$

The minimum of $D_K(P; Q)$ is obtained iff $p_i = q_i$ (see, e.g. [14]).

Apart of the Kullback-Leibler statistical divergence, a number of other divergence measures, depending on certain parameters, have been proposed and intensively studied by Rényi [5], Kapur [2], Kullback and Leibler [4], Havrda and Charvat [6], Tsallis [7], [8]. Some of them satisfy the convexity condition only for restricted values of the corresponding parameters.

*Electronic address: fyzimmar(at)savba.sk

†Electronic address: serge_shp(at)yahoo.com

However, the Kullback-Leibler, Rényi, Havrda-Charvat, Tsallis and the trigonometrical [9] statistical divergences require generally $p_i = 0$ whenever $q_i = 0$. From point of view of their application, this is not a desirable property because just such situations we often encounter in the theoretical physics especially in statistical and quantum physics.

In the next sections, we exploit one of the first statistical divergence measure, that was proposed in the literature, the Bhattacharyya divergence of P and Q which is symmetrical regarding the exchange of P and Q and does not suffer from the above-mentioned shortcoming. We attempt to apply the Bhattacharyya statistical divergence to the quantification of the degree of similarity of two quantum observables.

II. THE BHATTACHARYYA STATISTICAL DIVERGENCE

The Bhattacharyya statistical divergence of the discrete probability distributions

$$P = p_1, p_2, \dots, p_n; \quad Q = q_1, q_2, \dots, q_n$$

is defined as [10]

$$S(P, Q) = \sum_{i=1}^n (p_i q_i)^{1/2}. \quad (1)$$

This divergence has the following properties:

- (i) It becomes its maximal value equal to 1 when the probability distributions P and Q are identical.
- (ii) Its minimal value is zero when the components of P and Q do not overlap.
- (iii) Its value lies in the interval $[0, 1]$ and expresses the degree how much the probability distributions of P and Q are similar.
- (iv) It is *symmetrical* regarding the exchange of P and Q .
- (v) $S(P, Q)$ satisfies the properties of nonnegativity, finiteness and boundedness.
- (vi) It can be straightforward extended for more than two probability distributions [15].

The Bhattacharyya statistical divergence of P and Q has a simple geometrical interpretation. Consider the following vectors $\hat{P} \equiv \{\sqrt{p_1}, \sqrt{p_2}, \dots, \sqrt{p_n}\}$ and $\hat{Q} \equiv \{\sqrt{q_1}, \sqrt{q_2}, \dots, \sqrt{q_n}\}$. According to Eq. (1), the similarity measure of $\hat{P}$ and $\hat{Q}$ is simply the scalar product of $\hat{P}$ and $\hat{Q}$ in $\mathbf{R}_+^{(n)}$. Since $\hat{P}$ and $\hat{Q}$ represent the unit vectors in $\mathbf{R}_+^{(n)}$ its scalar product is equal to the cosine of angle between $\hat{P}$ and $\hat{Q}$ which, of course, has the properties (i)-(iv).

III. APPLICATION OF THE BHATTACHARYYA STATISTICAL DIVERGENCE TO QUANTUM MECHANICS

Consider two observables A, B with Hermitian operators $\hat{A}, \hat{B}$ in an N -dimensional Hilbert space, whose corresponding complete orthonormal sets of eigenvectors $\{|x_i^{(A)}\rangle\}, \{|x_i^{(B)}\rangle\}, \dots$ ($i = 1, 2, \dots, N$) are disjoint and have nondegenerate spectra. Let $|\phi\rangle$ be a normalized state vector of N -dimensional Hilbert space then it holds

$$|\phi\rangle = \sum_i^N a_i |x_i^{(A)}\rangle, \quad |\phi\rangle = \sum_j^N b_j |x_j^{(B)}\rangle, \dots$$

Accordingly, the components of the probability distributions $P(A)$ and $P(B)$ associated with the observables A and B are

$$P_i(A) = |a_i|^2 = |\langle x_i^{(A)} | \phi \rangle|^2; \quad \sum_{i=1}^n P_i(A) = \sum_{i=1}^n |a_i|^2 = 1 \quad (2a)$$

$$P_i(B) = |b_i|^2 = |\langle x_i^{(B)} | \phi \rangle|^2; \quad \sum_{i=1}^n P_i(B) = \sum_{i=1}^n |b_i|^2 = 1. \quad (2b)$$

Inserting Eqs.(2a) and (2b) into Eq.(1) we get

$$S(P(A), P(B)) = \sum_{i=1}^n (P_i(A) P_i(B))^{1/2} = \sum_{i=1}^n |a_i| |b_i|.$$

If $A = B$ then $|a_i| = |b_i|$ for $i = 1, 2, \dots, n$ and Eqs.(2a) and (2b) becomes

$$S(P(A), P(B)) = \sum_{i=1}^n P_i(a) = \sum_{i=1}^n |a_i|^2 = \sum_{i=1}^n P_i(B) = \sum_{i=1}^n |b_i|^2 = 1.$$

Hence, for $A \equiv B$ it follows $S(P(A), P(B)) = 1$. Given the state vector $|\phi\rangle$ and operators $\hat{A}, \hat{B}$ the considered statistical divergence of their probability distributions can be generally determined. To each operator, a ray in the Hilbert space can be assigned. The quantity $S(P(A), P(B))$ gives the closeness of different rays in Hilbert space. If these rays are identical then their Bhattacharyya divergence $S(P(A), P(B))$ is equal to 1. If they are perpendicular to each other then $S(P(A), P(B))$ becomes zero. Generally, $S(P(A), P(B))$ $A \equiv \hat{B}$ and the corresponding rays of these operators in Hilbert space are identical, i.e. the cosine of angle between them is equal to 1. Therefore, $S(P(A), P(B)) = 1$.

Next, we consider the case of two *non-commuting* observables. Consider two observables A and B with non-commuting Hermitian operators $\hat{A}$ and $\hat{B}$ in an N -dimensional Hilbert space, whose corresponding complete orthonormal sets of eigenvectors $\{|x_i\rangle\}$, $\{|y_i\rangle\}$ ($i = 1, 2, \dots, N$) are disjointed and have nondegenerate spectra. Let $|\phi\rangle$ be a normalized state vector of N -dimensional Hilbert space then it holds

$$|\phi\rangle = \sum_i^N a_i |x_i\rangle, \quad |\phi\rangle = \sum_j^N b_j |y_j\rangle.$$

According the quantum transformation theory we have

$$|\phi\rangle = \left(\sum_i^N a_i \langle x_i | y_1 \rangle \right) |y_1\rangle + \left(\sum_i^N a_i \langle x_i | y_2 \rangle \right) |y_2\rangle + \dots = \sum_j^N \sum_i^N a_i \langle x_i | y_j \rangle |y_j\rangle$$

$$P_i(A) = |\langle x_i | \phi \rangle|^2 = |a_i|^2, \quad Q_j(B) = |b_j|^2 = |\langle y_j | \phi \rangle|^2 = \left| \left(\sum_i^N a_i \langle y_i | x_j \rangle \right) \right|^2, \quad (3)$$

where $\langle x_i | y_j \rangle$ $i, j = 1, 2, 3, \dots, N$ are the elements of the transformation matrix $\mathbf{T}$ between the observables A and B

$$\mathbf{T} = \begin{pmatrix} \langle x_1 | y_1 \rangle & \langle x_1 | y_2 \rangle & \dots & \langle x_1 | y_n \rangle \\ & \dots & & \\ \langle x_n | y_1 \rangle & \dots & & \langle x_n | y_n \rangle \end{pmatrix}$$

Inserting Eq.(3) into Eq.(1) we get for the Bhattacharyya statistical divergence of the probability distributions $P(A)$ and $Q(B)$ the expression

$$S(P(A), Q(B)) = \sum_{i=1}^n \sqrt{P_i(A) Q_i(B)} = \sum_{i=1}^n |a_i| |b_i| = \sum_{i=1}^n |\langle x_i | \phi \rangle| |\langle y_i | \phi \rangle| = \sum_{j=1}^n |a_j| \left| \left(\sum_{i=1}^n a_i \langle y_i | x_j \rangle \right) \right|. \quad (1)$$

Given the state vector $|\phi\rangle$ and the components of $\mathbf{T}$, the divergence of the probability distributions of A and B can be generally determined.

Next, we present an example for determining of Bhattacharyya divergence of two concrete *complementary* observables describing a simple quantum system.

IV. AN EXAMPLE

For the sake of simplicity, we will consider the complementary observables in a two-dimensional Hilbert space. Such system represents a particle with spin $\hbar/2$ [13]. Determining the probability distributions of the components J_x and J_z we can calculate the similarity measure of their probability distributions.

The state vector of this quantum system is spinor

$$|\Psi\rangle = \begin{pmatrix} a_1 \\ a_2 \end{pmatrix},$$

where

$$a_1 a_1^* + a_2 a_2^* = 1.$$

Its wave functions in z-representation takes the form $|\Psi\rangle_z = a_1|z_1\rangle + a_2|z_2\rangle$. According Eq.(3), the probability P_{z_1} and P_{z_2} that J_z is projected on $|z_1\rangle$ and $|z_2\rangle$ is $|a_1|^2 = a_1 a_1^*$ and $|a_2|^2 = a_2 a_2^*$, respectively, so the corresponding probabilistic schema becomes

$$\frac{J_z}{P_{J_z}} \left\| \begin{array}{c} |z_1\rangle \\ |z_2\rangle \end{array} \right\| \begin{array}{c} a_1 a_1^* \\ a_2 a_2^* \end{array}$$

Using the transfer transformation matrix

$$\mathbf{T} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}$$

we obtain Ψ in its x-representation $|\Psi\rangle_x = \frac{a_1 + a_2}{\sqrt{2}} |x_1\rangle + \frac{a_1 - a_2}{\sqrt{2}} |x_2\rangle$. Similarly, the probabilistic scheme for J_x turns out to be

$$\frac{J_x}{P_{J_x}} \left\| \begin{array}{c} |x_1\rangle \\ |x_2\rangle \end{array} \right\| \begin{array}{c} 2^{-1}(a_1 + a_2)(a_1^* + a_2^*) \\ 2^{-1}(a_1 - a_2)(a_1^* - a_2^*) \end{array}.$$

Now, we express a_1 and a_2 by means of new variables r and φ in the following way

$$a_1 = \sqrt{r} \exp(i\varphi_1), \quad a_2 = (\sqrt{1-r}) \exp(i\varphi_2), \quad \varphi = \varphi_1 - \varphi_2.$$

In these variables, we obtain for J_x and J_z the following probability distributions

$$Q(J_x) \equiv \{r^2, (1-r)^2\}$$

and

$$P(J_z) \equiv \left\{ \frac{1}{2}(1 + 2\sqrt{r-r^2} \cos \varphi), \frac{1}{2}(1 - 2\sqrt{r-r^2} \cos \varphi) \right\}.$$

The similarity measure of these probability distributions consists of two terms

$$S(Q(J_x), P(J_z)) = T_1 + T_2, \quad (4)$$

where

$$T_1 = \sqrt{\frac{r}{2}(1 + 2\sqrt{r-r^2} \cos \varphi)} \text{ and } T_2 = \sqrt{\frac{(1-r)}{2}(1 - 2\sqrt{r-r^2} \cos \varphi)}.$$

If J_x occurs in one of its eigenstates, i.e. $r = 1$ or 0 , then the first or second term in Eq.(4) becomes zero and we obtain the minimal value of $S(P(J_x), Q(J_z))$ equal to $T_1 = T_2 = \sqrt{1/2}$. $S(P(J_x), Q(J_z))$ attains its maximal value equal to 1 for $r = \sqrt{1/2}$ and $\varphi = \pi/2$. The 3D-plot $S(P(J_x), Q(J_z))$ is given in Fig. 1.

This graph shows that in the vicinity of $r = 0.85$ and for $\varphi \in [0, \pi/2]$ a hump occurs, where the probability distributions are almost similar. $S(P(J_x), Q(J_z))$ never drops under the value $\sqrt{1/2}$, therefore it holds

$$S(P(J_x), Q(J_z)) \geq \sqrt{1/2}.$$

While $S(P:Q)$ for two commuting observables in two-dimensional Hilbert space can attain arbitrary value, the similarity measure of the probability distributions of the investigating non-commuting observables, J_x and J_z , is bounded by the value $\sqrt{1/2}$. We note that two observables A, B in an N -dimensional Hilbert space are said to be complementary (to each other) if their transformation matrix has the form [11]

$$|\langle a_i | b_j \rangle| = N^{-1/2} \quad (i, j = 1, \dots, N).$$

Complementary observables can be considered as a generalization to higher dimensions of spin-1/2 orthogonal system [12]. Hence, we can proceed similarly also for two complementary observables in a general N -dimensional Hilbert space.

The concept of Bhattacharyya statistical divergence is quite independent of quantum mechanics and can be defined in any probability space. Hence, apart from the application of the similarity measure in quantum physics it can also be applied to theory of the physical measurement. It may serve as a certain degree of the reliance of a physical measurement. Suppose that in two different laboratories the probabilities of the decay modes of an elementary particle are measured. Generally, the different probability distributions of the individual decay modes are obtained in each laboratory. To determine the reliance of the measurement we can insert the measured probability distributions in formula (1). Here, the simple rule holds: the large is the value of $S(P_1, P_2)$ the more reliable is the corresponding measurement.




FIG. 1: 3D-plot of the similarity measure of the probabilities $\mathbf{P}_{\mathbf{J}_x}$ and $\mathbf{P}_{\mathbf{J}_z}$ as function of r and φ .

Acknowledgments

Partial support by the Grant Agency VEGA No. 2/6087/26 is highly acknowledged.

-
- [1] I. Vajda, *Theory of statistical Inference and Information*. Dordrecht: Kluwer, 185.
 - [2] J. W. Karpur, J. Math. Phys. Sci. **17**, (1983), 103.
 - [3] S. Kullback, *Information Theory and Statistics*. (John Willey: New York) 1959.
 - [4] S. Kullback, R. A. Leibler, Ann. Math. Stat. **22**, (1951), 79. (1951).
 - [5] A. Rényi, "On the Measures of Entropy and Information". In: *4th Berkeley Symp. Math. Stat. Probability*, **1** (1961), 541.
 - [6] J. H. Havrda and F. Charvat, Kybernetika **3** (1967), 30.
 - [7] C. Tsallis, J. Stat. Phys. **52** (1988), 479.
 - [8] C. Tsallis, Brazilian J. Phys. **29** (1999), 1-87.
 - [9] V. Majerník, *Elementary theory of organization*. (Palacký University Press, Olomouc: 2001).
 - [10] A. Bhattacharyya, *On a Measure of Divergence Between Two Statistical Populations Defined by their Probability Distributions*, Bull. Calcutta Math. Soc. **35**. (1943), 99-109.
 - [11] K. Kraus, Phys. Rev. **D 35** (1987), 3070.
 - [12] J. Sánchez, Phys. Lett. **A 173** (1993), 233.
 - [13] E. Merzbacher, *Quantum Mechanics*, 2nd Ed. (Wiley, New York, 1972).
 - [14] G. Aczél and Dároczy, *On Measures of Information and Their Characterization*, Academic Press, New York, 1975.
 - [15] V. Majerník, Intern. J. Gen. Systems **33** (2004), 673.